

Clinical research of intraoperative image-guidance in endoscopic nasocular operation

Xiao-Yan Dou¹, Jiang Guo¹, Yong-Tian Lu², Jing Sima¹

Foundation item: Science Research Program of Shenzhen, China (No. 200802052)

¹Department of Ophthalmology, the Second People's Hospital of Shenzhen, Shenzhen 518035, Guangdong Province, China

²Department of Ear-nose-throat, the Second People's Hospital of Shenzhen, Shenzhen 518035, Guangdong Province, China

Correspondence to: Jing Sima. Department of Ophthalmology, the Second People's Hospital of Shenzhen, Shenzhen 518035, Guangdong Province, China. sima618@163.com

Received: 2010-06-10 Accepted: 2010-07-20

Abstract

• **AIM:** To evaluate the availability and our experience of intraoperative image-guidance in endoscopic nasocular operation.

• **METHODS:** Seven cases of endoscopic nasal surgery with intraoperative image-guidance were retrospectively reviewed, including 3 cases of optic nerve injury; 3 cases of foreign object of optic behind the eyeball; 1 case of retrobulbar tumor (angioma).

• **RESULTS:** The preoperative preparatory time would take 15 minutes, including coordination, head holder localization, conventional instrument registration. In our cases, the localization accuracy between 3-D image landmarks of navigation system and actual anatomical landmarks was less than 1.3mm. The optic nerve and other anatomical points could be orientated accurately in intraoperative procedures. No complication occurred.

• **CONCLUSION:** Nasal endoscope combined with image-guidance systems provides accurate anatomical localization of anterior skull base with enlarged operation field. It is possible for surgeons to observe important anatomical structures during endoscopic surgery. It could increase the effectiveness and decrease surgical complications, especially in complicated cases.

• **KEYWORDS:** operation on orbit; image-guidance systems; nasal endoscope

DOI: 10.3969/j.issn.1672-5123.2010.08.004

Dou XY, Guo J, Lu YT, Sima J. Clinical research of intraoperative image-guidance in endoscopic nasocular operation. *Int J Ophthalmol (Guoji Yanke Zazhi)* 2010;10(8):1467-1468

INTRODUCTION

Because the orbital complex, narrow space, a number of vital organs, so anatomical localization is very important for orbital surgery. Recently, technology for computer-assisted surgery was introduced and popularized to improve the overall safety of orbital surgery. This technology allows a direct

comparison of intraoperative anatomy with preoperative image information. In this study, a Medtronic, Inc. navigation System for navigation was used during transnasal endoscopic surgeries.

MATERIALS AND METHODS

Materials Since August 2008 to December 2009, 7 cases of endoscopic nasal surgery with intraoperative image-guidance were retrospectively reviewed, including 3 cases of optic nerve injury; 3 cases of foreign object of optic behind the eyeball; 1 case of retrobulbar tumor (angioma). Six males, 1 female, aged 23 to 61 years old, mean age was 45 years old.

Methods

Equipment Image-guided system: Medtronic, Inc.; Nasal endoscope: STOZE Inc., Germany; Power system operation: Medtronic XPS3000, Inc.

Preparing for surgery A high-resolution CT spiral scan with serial 1mm thick slices on a horizontal plane was obtained 1 or 2 days before surgery. The image data was transferred to a computer workstation via an optical disk to generate 3-D data, from which nine to eleven reference markers were selected for calibration before the surgical procedure.

At the beginning of each case, a special headset was placed on the patient's head, which permitted attachment of a reference flame with light-emitting diodes for optical tracking. The navigation functions were activated immediately after the position of the head was determined. The position of the tip of the inhaler was displayed on the workstation's screen indicating the location of the target and its adjacent normal structures, confirming the proposed operative field. In all cases, transnasal endoscopic surgery was performed as done previously.

RESULTS

Seven cases were successfully performed surgery without complications. Imaging signs and physical anatomical landmark error ≤ 1.5 mm. Optic nerve injury in 3 patients, 2 patients with preoperative visual acuity: no light perception and postoperative visual acuity: light perception. One patient with preoperative visual acuity: 0.1 and postoperative visual acuity: 0.5. Three cases of foreign body in orbit: foreign body was successfully removed no significant changes in visual acuity. One case of tumor resection, prognosis is good.

DISCUSSION

Because of the complexity of eyeball, orbit and sinuses, eye trauma often appeared with head trauma. Orbital surgery through nasal cavity has many advantages: more efficient, less damage and faster recovery. Application and development of nasal endoscopy promoted the rapid development of orbital

surgery^[1,2].

Because the optic nerve is very delicate, and endoscopic surgery is two-dimensional image, so image guidance for surgery is very important.

In this study we used Medtronic Inc. Stealth Station TM system. Image-guided system is to use electromagnetic or infrared signals to determine the precise location in surgery. System uses special computer software, the CT of MRI images of patients before surgery reconstruction, and through electromagnetic induction position on the operation instruments for precise positioning. Doctors operated under the guidance of three-dimensional images.

It is important to calibrate before surgery which has an effect on the accuracy of the navigation system. Medtronic Inc Stealth Station TM system demanded a one to one match of reference marks and points in the three-dimensional reconstruction image. In all cases of present study, the average time for preparation is 15 minutes.

Because the optic nerve in optic foramen is limited, and prone to fracture, optic nerve injury is a kind of so common traumatic eye disease^[3]. Current traumatic optic neuropathy treatment is still much controversy, but most scholars believe that surgical treatment can improve the efficiency in a clear imaging diagnosis of optic canal fracture, cavity deformation, hematoma compression or early high-dose hormone treatment fails.

Li and Xu^[4] found that surgical treatment of traumatic optic neuropathy is important. Transnasal endoscopic optic canal decompression is commonly used in clinical surgery. In recent years, a lot of successful cases at home and abroad have been reported, with fewer traumas, clear vision and less scar^[5,6].

Image-guided system further reduces surgical risk, increase the success rate. Optic nerve trauma are usually associated with compound fractures, loss of normal anatomic marks^[7], endoscopic surgery can not accurately determine the local anatomy. Image-guided system can help doctors determine the exact anatomical location, provide accurate three-dimensional images. It reduces serious complications, improve surgical accuracy and safety.

Patients with optic nerve injury accept VEP examination to evaluate visual function and to judge results of operations, some scholars have reported that accuracy was 90%^[8]. But VEP is not the criteria to decide whether surgery, even if it results very poor, we believe that should not give up easily. When doctors found fractures, fracture fragments on the optic nerve compression or disc edema, the patient should undergo surgery as soon as possible. Accurate identification of the optic canal is the key to successful operation^[9]. After surgery, although visual acuity may be better, but optic nerve atrophy is still possible.

Orbital foreign body can lead to a variety of dysfunction. It can affect eye movement, diplopia occur and affect the optic nerve blood supply, leading to optic atrophy^[3]. Because the orbital space narrow, and the orbital tissue loose, so doctors are easy to slip foreign body, likely to cause tissue damage. With system's help, doctors can avoid the optic nerve and other vital organs, operate safely and quickly.

There are some problems associated with the clinic application of the navigation system; Preoperative image examination can not reflect changes in the operation completely; A loose headset can distort the correspondence of one to one matching of reference points and dislocate the structure during surgery; The navigation system required more expenses for instruments and increased the cost of an in-patient. Therefore, doctors need to evaluate comprehensively, select treatment options, in order to achieve better clinical results.

REFERENCES

- 1 Yang ZQ. The development of rhino-neurosurgery and rhino-eyesurgery. *Chin J Otorhinolaryngol* 1999;34(3):273-275
- 2 Xv G, Li Y. Jinan University Press. Endoscopic sinus surgery 1994; 120-122
- 3 Li FM, Eye Book. People's Medical Publishing House 1996;3441-3442
- 4 Li Y, Xu G. Traumatic optic neuropathy: Importance and result of surgical treatment. *Chin J Otorhinolaryngol* 2002;37(4):206-209
- 5 Kountakis SE, Maillard AA, El-Harazi SM, Longhini L, Urso RG. Endoscopic optic nerve decompression for traumatic blindness. *Otolaryngol Head Neck Surg* 2000;123 (1 Pt 1):34-37
- 6 Luxenberger W, Stammberger H, Jeebles JA, Walch C. Endoscopic optic nerve decompression; the Graz experience. *Laryngoscope* 1998;108 (6):873-882
- 7 Ge WT, Zhou B, Han DM. Evaluation of imaging navigation system during endoscopic sinus surgery. *Chin Arch Otolaryngol Head Neck Surg* 2006;13(8):565-569
- 8 Wang ZQ, Ma ZZ, Jiang L. The clinical value of Visual Evoked Potential in traumatic optic neuropathy. *Chinese Ophthalmic Research* 2002;20(3):261-263
- 9 Sun Y, Li N, Yang SK. Otorhinolaryngology Head and Neck Surgery Techniques. Beijing Science Press 2004:144-149

影像导航在鼻内镜下眼眶手术的临床研究

窦晓燕¹, 郭 疆¹, 卢永田², 司马晶¹

基金项目:中国广东省深圳市科技计划资助项目(No. 200802052)
(作者单位:518035 中国广东省深圳市第二人民医院暨深圳大学第一附属医院¹眼科;²耳鼻喉科)

作者简介:窦晓燕,女,主任医师,研究方向:眼屈光手术及眼外伤的手术治疗。

通讯作者:司马晶,女,教授,主任医师,硕士研究生导师,2001/2003年在美国南卡罗纳医科大学眼科从事博士后研究。现任广东省视光学会视力保健专业委员会副主任委员,深圳市医学会眼科专业委员会副主任委员等职务,研究方向:白内障、小儿眼科及眼视光学. sima618@163.com

摘要

目的:探讨影像导航技术在鼻内镜眼眶微创手术中应用。

方法:回顾性分析7例患者,其中视神经损伤3例;眶内球后异物3例;球后肿瘤1例(血管瘤)。所有病例均采用Brain Lab影像导航系统及鼻内镜实施手术。

结果:影像导航术前准备时间(包括配准、头架定位、常规器械注册等)10~20(平均15)min。手术区域影像标志与实体解剖标志间的误差 ≤ 1.3 mm。影像导航可准确定位视神经等解剖标志,无手术并发症发生。

结论:影像导航系统可以帮助术者在术中定位眼眶内重要的解剖结构,并可使手术视野扩展到内镜之外,使术者在术野中进行手术操作的同时,能顾及到术野周围的重要结构。特别是在复杂的手术中,可提高手术的精确性和安全性。

关键词:眼眶手术;影像导航;鼻内镜