

# Effect of Tetramethylpyrazine on RPE degeneration, choroidal blood flow and oxidative stress of RPE cells

Yi Shen, Pei Zhuang, Bao-Qin Lin, Wan-Yu Zhang, George C Y Chiou

Institute of Ocular Pharmacology, College of Medicine, Texas A&M Health Science Center, College Station, TX 77843, USA

**Correspondence to:** George C Y Chiou. Institute of Ocular Pharmacology, College of Medicine, Texas A&M Health Science Center, College Station, TX 77843, USA. chiou@medicine.tamhsc.edu

Received:2010-06-28 Accepted:2010-08-23

## Abstract

• **AIM:** To study the effects of Tetramethylpyrazine (TMP) on retinal pigment epithelium (RPE) degeneration, choroidal blood flow and oxidative stress of RPE cells.

• **METHODS:** The 35mg/kg NaIO<sub>3</sub>-induced RPE degeneration rat eyes was given 25μg 1% TMP eye drops 3 times a day for 7 days before NaIO<sub>3</sub> injection, and then 2 to 4 weeks after NaIO<sub>3</sub> injection. RPE function was measured with c-wave of electroretinogram (ERG). Colored microsphere technique was used for *in vivo* experiments to determine the choroidal blood flow in ocular hypertensive (40mmHg) rabbit eyes. Methylthiazolotetrazolium (MTT) assay was used to study *in vitro* effect of TMP on various oxidants induced injury in the hRPE (ARPE-19 (ATCC, Manassas, VA, USA)).

• **RESULTS:** Two weeks after NaIO<sub>3</sub> injection, the amplitude of ERG c-wave fell markedly in NaIO<sub>3</sub> group to 36% of control group ( $P < 0.01$ ). No apparent difference was observed in TMP + NaIO<sub>3</sub> group. Four weeks later, the NaIO<sub>3</sub> group fell to 46% of control group ( $P < 0.01$ ), while the TMP + NaIO<sub>3</sub> group fell to only 77% of control group ( $P < 0.01$ ). There was a 67% reversal of the ERG c-wave by TMP as compared to NaIO<sub>3</sub> group ( $P < 0.01$ ). The choroidal blood flow was significantly increased at all time points (at 30, 60 and 120 minutes after TMP instillation) as compared with corresponding controls. TMP had no effect on hypoxia-(1% O<sub>2</sub>), t-BHP- and H<sub>2</sub>O<sub>2</sub>-induced damage in RPE cells. 10μg/mL TMP could reverse 1 and 3mmol/L NaN<sub>3</sub>-induced loss of viability of RPE by 18.5% ( $P < 0.01$ ) and 23% ( $P < 0.01$ ), respectively. 30μg/mL TMP could reverse 30 and 100mmol/L NaIO<sub>3</sub> induced loss of viability of RPE by 18.1% ( $P < 0.05$ ) and 16.8% ( $P < 0.01$ ), respectively.

• **CONCLUSION:** TMP can significantly protect RPE from NaIO<sub>3</sub> induced degeneration *in vivo* and oxidative stress *in vitro* and can increase choroidal blood flow markedly *in vivo*.

• **KEYWORDS:** Tetramethylpyrazine; sodium iodate; retinal pigment epithelium; age-related macular

degeneration; choroidal blood flow; oxidative stress  
DOI:10.3969/j.issn.1672-5123.2010.10.001

Shen Y, Zhuang P, Lin BQ, Zhang WY, Chiou GCY. Effect of Tetramethylpyrazine on RPE degeneration, choroidal blood flow and oxidative stress of RPE cells. *Int J Ophthalmol(Guji Yanke Zazhi)* 2010;10(10):1843-1887

## INTRODUCTION

A 2004 analysis reported that among Americans over the age of 40, AMD and/or geographic atrophy were present in at least one eye in 1.47% of the population, and that 1.75 million individuals have AMD. The prevalence of AMD increased dramatically with age, with more than 15% of the white women older than 80 years having neovascular AMD and/or geographic atrophy. More than 7 million individuals are at substantial risk of developing AMD. Owing to the rapidly aging population, the number of persons having AMD will increase by 50% to 2.95 million in 2020<sup>[1]</sup>. In another study, AMD was reported to account for 54% of all current cases of blindness among the Caucasian population in the United States<sup>[2]</sup>. The study predicted that the number of blind people in the US could increase by as much as 70% by 2020. Traditional Chinese Medicine, Ligusticum Wallichii Franchet (Chuan Xiong), has been used to treat neurovascular and cardiovascular diseases. Tetramethylpyrazine (TMP) is one of the most active ingredients of Chuan Xiong. TMP can increase retinal and choroidal blood flow by 44% and did not affect systemic blood pressure and heart rate on rabbit at the same time<sup>[3]</sup>. Intraperitoneal administration of TMP on Brown Norway rats could significantly decrease the intensity of fluorescein leakage from the photocoagulated lesions and the size of choroidal neovascularization (CNV) induced by laser treatment<sup>[4]</sup>. TMP could protect the heart from lipid peroxidation-induced heart toxicity by its prominent anti-lipid peroxidation and anti-free radical formation effects<sup>[5]</sup>. This study is to observe the effects of TMP on sodium iodate (NaIO<sub>3</sub>) induced retinal pigment epithelium (RPE) degeneration, choroidal blood flow *in vivo* and oxidative stress of RPE cells *in vitro*.

## MATERIALS AND METHODS

**Materials** TMP (purity ≥98%), NaIO<sub>3</sub> (purity ≥99.5%), thiazolyl blue tetrazolium bromide (MTT, purity ≥97.5%), hydrogen peroxide (50wt% solution in water), tert-Butyl hydroperoxide (t-BHP, 70wt% in water), dimethyl sulfoxide (DMSO, purity ≥99.9%), sodium azide (purity ≥99.5%), and Dulbecco's modified Eagles's medium/Ham's F12

(DMEM/F12, 1:1) were purchased from Sigma-Aldrich Chemical Co. (St. Louis, MO, USA). Fetal bovine serum (FBS) was purchased from GIBCO (Grand Island, NY, USA).

## Methods

**ERG in rats** A total of thirty 8-week-old male Brown-Norway (BN) rats were randomly divided into 3 groups, 10 rats in normal group, 10 rats in NaIO<sub>3</sub> group and 10 rats in TMP + NaIO<sub>3</sub> group. Control group was instilled with solvent (2-hydroxypropyl-β-cyclodextrin, Sigma-Aldrich) alone without NaIO<sub>3</sub> injection. NaIO<sub>3</sub> group was instilled with solvent plus 35mg/kg NaIO<sub>3</sub> injection through hypoglossal vein, whereas TMP + NaIO<sub>3</sub> group was instilled with 1% TMP eye drops plus 35mg/kg NaIO<sub>3</sub> injection<sup>[6,7]</sup>. Both eyes of all rats were instilled with 1 drop for 3 times a day for 1 week before and 4 weeks after NaIO<sub>3</sub> injection<sup>[8]</sup>. At the end of 2 and 4 weeks, RPE function was measured with c-wave of ERG.

BN rats were dark adapted for 2 hours, and then anesthetized with ketamine 35mg/kg plus xylazine 5mg/kg *i. m.* Half of the initial dose was given each 1 hour thereafter. Pupils of all rats were dilated with one drop each of 1% atropine, and 2.5% phenylephrine. Before recording, one drop of 0.5% tetracaine was given for surface anesthesia. All rats were kept warm during ERG measurement. DC-ERG recording methods developed by Peachey were followed. Briefly, a 1-mm diameter glass capillary tube with filament (Sutter Instruments, Novato, CA) that was filled with Hanks balanced salt solution (Invitrogen, Carlsbad, CA) was used to connect with a Ag/AgCl wire electrode with an attached connector. The electrode was positioned to the corneal surface. Responses were amplified (dc-100 Hz; gain = 1000X; DP-31, Warner Instruments, Hamden, CT) and digitized at 10Hz or 1000Hz. Data were analyzed by iWORX LabScribe Data Recording Software (iWorx0CB Sciences, Dover, NH). Light stimuli was derived from an optical channel using a fiber-lite high intensity illuminator (Dolan-Jenner Industries, Inc. MA), with neutral density filters (Oriel, Stratford, CT) placed in the light path to adjust stimulus luminance. The stimulus luminance used in this experiment was 3.22 log cd/m<sup>2</sup> and stimulated for 4 minutes. Luminance calibration was made by a Minolta (Ramsey, NJ) LS-110 photometer focused on the output side of the fiber optic bundle where the rat eye was located.

## Measurement of Choroid Blood Flow in Ocular Hypertensive Rabbit Eyes

New Zealand white rabbits, weighing 2.5-3.0kg, were purchased through LARR (Texas A&M University, USA). All rabbits were anesthetized with 35mg/kg ketamine and 5mg/kg xylazine *i. m.* Half of the initial dose was given each 1 hour thereafter. An ocular hypertensive model was created by raising the intraocular pressure of the left eye to 40mmHg, which reduced the ocular blood flow to approximately one third of the normal value<sup>[9]</sup>. The left ventricle was cannulated through the right carotid artery for the injection of colored microspheres and the femoral artery was cannulated for blood sampling. One percent drug solution (50μL) or vehicle (50μL) was instilled topically to the left eye, and the ocular blood flow of the ocular

hypertensive rabbits was measured with colored microspheres at 0, 30, 60 and 120 minutes thereafter. At each time point, 2 million microspheres in 0.2mL were injected as a reference, and blood samples were taken from the femoral artery for exactly 1 minute following injection of the microspheres. The blood sample was collected in a heparinized tube and the volume was recorded. The rabbits were euthanized with an injection of 100mg/kg pentobarbital sodium after the last blood sampling. The left eyes were enucleated and dissected into the retina, choroid, iris and ciliary body. The tissue samples were weighed.

The details of sample processing and microsphere counting were provided by E-Z Trac (Los Angeles, CA). In brief, a hemolysis reagent was added to the microfuge tubes with the blood sample, then vortexed and centrifuged for 30 minutes at 6000r/min. The supernatant was removed, and tissue/blood digest reagents I and II were added. The tubes were capped, vortexed, and centrifuged for 30 minutes again. The supernatant was removed, and the counting reagent was added, then vortexed, and centrifuged for 15 minutes at the same resolutions as above. The supernatant was removed, and the microspheres were resuspended in a precise volume of the counting reagent, the number of microspheres was counted with a hemocytometer.

Tissue/blood digest reagent I was added to the microfuge tubes with the tissue samples, sealed, and heated at 95°C for 15 minutes. The tubes were vortexed for 30 seconds, then reheated and revortexed until all tissue samples were dissolved. The tissue/blood digest reagent II was added while the tissue samples were still hot, then the tubes were capped, vortexed, and centrifuged for 30 minutes. The protocol thereafter was the same as that used to process the blood sampled, and the microspheres were counted.

The blood flow of each tissue at a certain time point was calculated from the following equation:  $Q_m = (C_m \times Q_r) / C_r$ , in which  $Q_m$  is the blood flow of a tissue in terms of μL/minute per mg,  $C_m$  is the microsphere count per mg of tissue,  $Q_r$  is the flow rate of blood sample in terms of μL per minute, and  $C_r$  is the total microsphere count in the referenced blood sample.

**Oxidative Stress of RPE Cells** The hRPE cells line, ARPE-19 (ATCC, Manassas, VA, USA), was used during all experiments. The cells were grown in DMEM/F12 supplemented with 10% FBS, 100units/mL penicillin G, and 100μg/mL streptomycin sulfate. Cells were incubated in a humidified incubator at 37°C under 5% CO<sub>2</sub> and 95% air.

The effect of TMP and oxidizing agents on the cell proliferation was evaluated with the MTT assay. RPE cells ( $8 \times 10^4$  cells) were seeded in 96-well plates (100μL/well) and allowed to grow overnight. Blanks were formed by adding 100μL medium (OD<sub>blank</sub>). The cells were then treated with fresh medium with TMP and/or oxidizing agents at the same time for 12, 24, or 48 hours (200μL/well, OD<sub>compound</sub>). The vehicle control group (OD<sub>control</sub>) was treated with 1.25% DMSO and/or PBS. The medium was then replaced with fresh medium containing 0.5mg/mL MTT but free FBS for 4 hours, 100μL/well. After

incubation, the medium was discarded and 150µL DMSO was added to solubilize formazan produced from MTT by the viable cells. Absorbance was measured at 570nm using a microplate reader (Packard BioScience Co. Meriden, CT, USA). Cell viability was calculated according to the following formula:

$$\text{Viability of cell (\%)} = (\text{OD}_{\text{compound}} - \text{OD}_{\text{blank}}) / (\text{OD}_{\text{control}} - \text{OD}_{\text{blank}}) \times 100\%$$

RPE cells were allowed to attach overnight, and then exposed to TMP and vehicle under hypoxic condition for 24 hours. Hypoxic conditions (1% O<sub>2</sub> and 5% CO<sub>2</sub>) were maintained by using a temperature- and humidity-controlled environmental C-chamber by O<sub>2</sub> and CO<sub>2</sub> controllers (Proox Model 110 and Pro CO<sub>2</sub> Model 120, Bio Spherix Ltd., Redfield, NY) with N<sub>2</sub> and CO<sub>2</sub> gas sources. Hypoxic culture medium was pre-equilibrated overnight prior to cell exposure.

**Statistical Analysis** All data were presented as mean ± SEM. A nonpaired Student's *t*-test was performed to analyze the significance between two means at a certain time point. The differences were considered significant at *P* < 0.05.

**RESULTS**


**ERG Recordings** ERG c-wave of rats was measured at 2 and 4 weeks after administration of 35mg/kg NaIO<sub>3</sub>. It was found that 2 weeks after NaIO<sub>3</sub> injection (Figure 1A), the amplitude of ERG c-wave of NaIO<sub>3</sub> group fell markedly to 36% of control group (*P* < 0.01). No significant change was observed in TMP + NaIO<sub>3</sub> group as compared to control. Four weeks after NaIO<sub>3</sub> administration (Figure 1B), the ERG c-wave of NaIO<sub>3</sub> group fell to 46% of control group (*P* < 0.01) whereas the TMP + NaIO<sub>3</sub> group only fell to 77% of control group (*P* < 0.01). There was a significant reversal of NaIO<sub>3</sub> group by TMP (Control: 0.393 ± 0.046 millivolts; NaIO<sub>3</sub>: 0.183 ± 0.058 millivolts; TMP + NaIO<sub>3</sub>: 0.305 ± 0.041 millivolts, *P* < 0.01).

**Choroid Blood Flow in Ocular Hypertensive Rabbit Eyes**


The choroid blood flow declined gradually in the vehicle control group. TMP significantly increased the choroid blood flow at time 30, 60 and 120 minutes as compared with the vehicle control group (Figure 2).

**Effect of TMP on Oxidants-Induced Damage in RPE Cells** TMP, up to the concentration of 30µg/mL, did not inhibit the proliferation of RPE cells (Figure 3). The result indicated that TMP itself had no toxicity on RPE cells at the concentration used in the experiments.


TMP has no effect on hypoxia, t-BHP, and H<sub>2</sub>O<sub>2</sub>-induced injury in RPE cells (data not shown). However, TMP reversed NaN<sub>3</sub>-induced injury in RPE cells (Figure 4B). 10µg/mL TMP reversed 1 and 3mmol/L NaN<sub>3</sub>-induced loss of viability of RPE by 18.5% (*P* < 0.01) and 23.0% (*P* < 0.01), respectively. 30µg/mL TMP reversed 0.3 and 3mmol/L NaN<sub>3</sub>-induced loss of viability of RPE from 73.8% to 91.0% (*P* < 0.01) and from 64.7% to 83.6% (*P* < 0.01), respectively. NaIO<sub>3</sub> concentration-dependently reduced the viability of RPE cells. 100µg/mL NaIO<sub>3</sub> significantly inhibited the proliferation of RPE cells (*P* < 0.05, Figure 5A). TMP significantly reversed NaIO<sub>3</sub>-induced injury in RPE cells (Figure 5B). 30µg/mL TMP reversed 30 and 100mmol/L NaIO<sub>3</sub>-induced loss



**Figure 1** A: Effect of TMP on NaIO<sub>3</sub> induced RPE degeneration in rat eyes for 2 weeks. a: control group compares with NaIO<sub>3</sub> group (*P* = 4.83E-9). b: NaIO<sub>3</sub> + 0.5% TMP group compares with NaIO<sub>3</sub> group (*P* = 9.78E-5); B: Effect of TMP on NaIO<sub>3</sub> induced RPE degeneration in rat eyes for 4 weeks. a: control group compares with NaIO<sub>3</sub> group (*P* = 8.98E-6). b: NaIO<sub>3</sub> + 0.5% TMP group compares with NaIO<sub>3</sub> group (*P* = 3.92E-5).




**Figure 2** Effect of TMP on the choroid blood flow of rabbit eyes. <sup>a</sup>*P* < 0.05 and <sup>b</sup>*P* < 0.01 vs vehicle control group. The data was expressed as mean ± SEM, in the vehicle control group *n* = 7 and in the TMP group *n* = 6.




**Figure 3** Effect of TMP on the proliferation of RPE cells. RPE cells were incubated with TMP for 48 hours.

of viability of RPE by 18.1% (*P* < 0.05) and 16.8% (*P* < 0.01), respectively. 100µg/mL TMP reversed 30 and 100mmol/L NaIO<sub>3</sub>-induced loss of viability of RPE from 80.0% to 97.3% (*P* < 0.05) and from 55.5% to 84.2% (*P* < 0.01), respectively.



**Figure 4 Effect of TMP on NaNO<sub>3</sub>-induced ischemia toxicity in RPE cells** A: Effect of NaNO<sub>3</sub> on the proliferation of RPE cells; B: Effect of TMP on NaNO<sub>3</sub>-induced ischemia toxicity in RPE cells. RPE cells were incubated with TMP and NaNO<sub>3</sub> for 48 hours. <sup>a</sup>*P* < 0.05 and <sup>b</sup>*P* < 0.01 vs vehicle control. <sup>d</sup>*P* < 0.01 vs model control. The data was expressed as mean ± SEM, *n* = 6.



**Figure 5 Effect of TMP on NaIO<sub>3</sub>-induced injury in RPE cells** A: Effect of NaIO<sub>3</sub> on the proliferation of RPE cells; B: Effect of TMP on NaIO<sub>3</sub>-induced injury in RPE cells. RPE cells were incubated with TMP and NaIO<sub>3</sub> for 48 hours. <sup>a</sup>*P* < 0.05 and <sup>b</sup>*P* < 0.01 vs vehicle control. <sup>c</sup>*P* < 0.05 and <sup>d</sup>*P* < 0.01 vs model control. The data was expressed as mean ± SEM, *n* = 6.

## DISCUSSION

AMD, is divided into wet-AMD and dry-AMD, and is a debilitating disease of the eye, which manifests clinically with loss of central vision and pathologically with the accumulation of drusen, RPE degeneration, photoreceptor atrophy, and in wet AMD cases, with CNV formation. There are several risk factors, including age, race, smoking, and diet<sup>[10]</sup>. But the etiology and pathogenesis of the disease remain largely unclear. For wet-AMD, there are a number of established treatment options, such as laser photocoagulation, intravitreal corticosteroids, verteporfin photodynamic therapy (V-PDT), anti-angiogenic factors, surgery, and combination of several treatments<sup>[11]</sup>. Most optional treatments are for the wet-AMD, not for dry-AMD. There is no effective treatment today for the most prevalent dry-AMD<sup>[12]</sup>.

Dry-AMD is triggered by abnormalities in RPE that lies beneath the photoreceptor cells and normally provides critical nutritional and metabolic support to these light-sensing cells. Secondary to RPE dysfunction, macular rods and cones degenerate leading to the irreversible loss of vision. Oxidative stress, formation of drusen, accumulation of lipofuscin, local inflammation and reactive gliosis represent the pathologic processes implicated in pathogenesis of dry-AMD. The direct toxic effect of NaIO<sub>3</sub> on RPE cells with secondary effects on photoreceptors and the choriocapillaries *in vivo* is well known<sup>[13]</sup>. The mechanisms of the toxicity of NaIO<sub>3</sub> to RPE cells are as follows: first, NaIO<sub>3</sub> can increase the ability of melanin to convert glycine into glyoxylate, a potential cell toxic compound<sup>[14]</sup>; second, NaIO<sub>3</sub> could denaturant retinal proteins by changes of SH levels in retina<sup>[15]</sup>; third, NaIO<sub>3</sub> could cause considerable structure changes by breakdown of RPE diffusion barrier or by reduction of adhesion between RPE and photoreceptor cells<sup>[16-19]</sup>; finally, NaIO<sub>3</sub> inhibits various enzyme activities, such as triose phosphate dehydrogenase, succinodehydrogenase and lactate dehydrogenase<sup>[18,20]</sup>.

The ERG results showed that TMP can reverse NaIO<sub>3</sub>-induced injury in RPE cells by 67% at the end of 4 weeks. TMP showed protective effect against NaIO<sub>3</sub>-induced RPE degeneration in rat eyes. The NaIO<sub>3</sub> intoxication causes death of RPE cells and photoreceptor damage followed by marked phagocytic activity of proliferating de-differentiated pigmental cells leading to the final pigmentary picture of the fundus. Further morphological study is needed to reveal the TMP role in RPE protection. Our previous study showed that TMP could significantly decrease the intensity of fluorescein leakage from the photocoagulated lesions and the size of CNV induced by laser treatment on Brown Norway rats, and interfered with vascular endothelial cell proliferation *in vitro*<sup>[4]</sup>.

The choroid blood flow results showed significant increase of choroid blood flow by TMP. The increase of choroid blood flow may facilitate removal of metabolic wastes and replenish nutrients to RPE and photoreceptors, thus it may change the microenvironment, change the balance between pro- and antiangiogenesis factors, and then the change of the process of angiogenesis.

Mitochondria are the powerhouse of the cell, and their primary

function is to generate ATP through oxidative phosphorylation via the electron transport chain<sup>[21]</sup>.  $\text{NaN}_3$  is an inhibitor of cytochrome oxidase and catalase, and will downregulate electron transport and  $\text{O}_2$  consumption to cause the death of cells<sup>[22]</sup>. TMP could reverse  $\text{NaN}_3$ -induced injury in RPE cells (Figure 4), but have no effect on hypoxia-induced damage (1%  $\text{O}_2$ ) in RPE cells. The results indicated that TMP can reverse  $\text{NaN}_3$ -induced ischemia by protecting cytochrome oxidase and/or catalase. Our experiments also showed that TMP reversed  $\text{NaIO}_3$ -induced injury in RPE cells *in vitro* (Figure 5).

In conclusion, TMP might slow the oxidative process of RPE cell layer which leads to RPE degeneration. TMP might also prevent the formation of CNV. According to the rational of the AMD, the RPE abnormalities or degeneration is the key point of both dry and wet AMD, so TMP could be used to prevent and treat both dry and wet AMD in the future.

#### REFERENCES

- 1 Friedman DS, O'Colmain BJ, Muñoz B, Tomany SC, McCarty C, de Jong PT, Nemesure B, Mitchell P, Kempen J; Eye Diseases Prevalence Research Group. Prevalence of age-related macular degeneration in the United States. *Arch Ophthalmol* 2004;122(4):564-572
- 2 Congdon N, O'Colmain B, Klaver CC, Klein R, Muñoz B, Friedman DS, Kempen J, Taylor HR, Mitchell P; Eye Diseases Prevalence Research Group. Causes and prevalence of visual impairment among adults in the United States. *Arch Ophthalmol* 2004;122(4):477-485
- 3 Chiou GC, Yan HY, Lei XL, Li BH, Shen ZF. Ocular and cardiovascular pharmacology of tetramethylpyrazine isolated from *Ligusticum wallichii* Franch. *Zhongguo Yaoli Xuebao* 1991;12(2):99-104
- 4 Zou YH, Jiang W, Chiou GC. Effect of tetramethylpyrazine on rat experimental choroidal neovascularization *in vivo* and endothelial cell cultures *in vitro*. *Curr Eye Res* 2007;32(1):71-75
- 5 Liu CF, Lin CH, Chen CF, Huang TC, Lin SC. Antioxidative effects of tetramethylpyrazine on acute ethanol-induced lipid peroxidation. *Am J Chin Med* 2005;33(6):981-988
- 6 Grignolo A, Orzalesi N, Calabria GA. Studies on the fine structure and the rhodopsin cycle of the rabbit retina in experimental degeneration induced by sodium iodate. *Exp Eye Res* 1966;5(1):86-97
- 7 Wei J, Chiou GC. Effects of hydralazine on  $\text{NaIO}_3$ -induced rat retinal pigment epithelium degeneration. *Int J Ophthalmol* 2008;8(8):1504-1510
- 8 Hughes PM, Olejnik O, Chang-Lin JE, Wilson CG. Topical and systemic drug delivery to the posterior segments. *Adv Drug Deliv Rev* 2005;57(14):2010-2032. Epub 2005 Nov 10
- 9 Chiou GC, Chen YJ. Effects of D- and L-isomers of timolol on retinal and choroidal blood flow in ocular hypertensive rabbit eyes. *J Ocul Pharmacol* 1992;8(3):183-190
- 10 Park YH, Chiou CY. Structure-activity relationship (SAR) between some natural flavonoids and ocular blood flow in the rabbit. *J Ocul Pharmacol Ther* 2004;20(1):35-42
- 11 Peachey NS, Stanton JB, Marmorstein AD. Noninvasive recording and response characteristics of the rat dc-electroretinogram. *Vis Neurosci* 2002;19(6):693-701
- 12 Coleman HR, Chan CC, Ferris FL 3rd, Chew EY. Age-related macular degeneration. *Lancet* 2008;372(9652):1835-1845
- 13 Kiuchi K, Yoshizawa K, Shikata N, Moriguchi K, Tsubura A. Morphologic characteristics of retinal degeneration induced by sodium iodate in mice. *Curr Eye Res* 2002;25(6):373-379
- 14 Baich A, Ziegler M. The effect of sodium iodate and melanin on the formation of glyoxylate. *Pigment Cell Res* 1992;5(6):394-395

- 15 Sorsby A, Reading HW. Experimental degeneration of the retina. XI. The effect of sodium iodate on retinal -SH levels. *Vis Res* 1964;4(10):511-514
- 16 Flage T, Ringvold A. The retinal pigment epithelium diffusion barrier in the rabbit eye after sodium iodate injection. A light and electron microscopic study using horseradish peroxidase as a tracer. *Exp Eye Res* 1982;34(6):933-940
- 17 Sen HA, Berkowitz BA, Ando N, de Juan E Jr. *In vivo* imaging of breakdown of the inner and outer blood-retinal barriers. *Invest Ophthalmol Vis Sci* 1992;33(13):3507-3512
- 18 Ashburn FS Jr, Pilkerton AR, Rao NA, Marak GE. The effects of iodate and iodoacetate on the retinal adhesion. *Invest Ophthalmol Vis Sci* 1980;19(12):1427-1432
- 19 Stern WH, Ernest JT, Steinberg RH, Miller SS. Interrelationships between the retinal pigment epithelium and the neurosensory retina. *Aust J Ophthalmol* 1980;8(4):281-288
- 20 Enzmann V, Row BW, Yamauchi Y, Kheirandish L, Gozal D, Kaplan HJ, McCall MA. Behavioral and anatomical abnormalities in a sodium iodate-induced model of retinal pigment epithelium degeneration. *Exp Eye Res* 2006;82(3):441-448. Epub 2005 Sep 19
- 21 Wang ZJ, Liang CL, Li GM, Yu CY, Yin M. Stearic acid protects primary cultured cortical neurons against oxidative stress. *Acta Pharmacol Sin* 2007;28(3):315-326
- 22 Thompson JG, McNaughton C, Gasparini B, McGowan LT, Tervit HR. Effect of inhibitors and uncouplers of oxidative phosphorylation during compaction and blastulation of bovine embryos cultured *in vitro*. *J Reprod Fertil* 2000;118(1):47-55

## Tetramethylpyrazine 在视网膜色素上皮细胞变性及脉络膜血流和 RPE 细胞氧化应激中的作用

沈 轶, 庄 沛, 林宝琴, 张婉玉, George C Y Chiou  
(作者单位: 美国德克萨斯州大学城, 德克萨斯农工大学医学院眼科药理学研究所)

通讯作者: George C Y Chiou. chiou@medicine.tamhsc.edu

### 摘要

**目的:** 研究 1% Tetramethylpyrazine (TMP) 在视网膜色素上皮细胞 (RPE) 变性, 脉络膜血流和 RPE 细胞氧化应激中的作用。

**方法:** 在碘酸钠诱导的大鼠 RPE 变性研究中, 1% TMP 滴眼液预先处理 1wk, 3 次/d, 1wk 后予碘酸钠舌下静脉注射, 在 2wk 和 4wk 末, 视网膜电图 (ERG) 测量 c 波。色素微球体技术分析高眼压状态下 TMP 对脉络膜血流的影响。Methylthiazoltetrazolium (MTT) 分析 TMP 在各种氧化应激中对 RPE 的保护作用。

**结果:** 碘酸钠注射后 2wk, 碘酸钠组 ERG 的 c 波下降至对照组的 36% ( $P < 0.01$ )。4wk 后, 碘酸钠组下降至对照组的 46% ( $P < 0.01$ ), 而 1% TMP + 碘酸钠组下降至对照组的 77% ( $P < 0.01$ )。与碘酸钠组比较, 1% TMP + 碘酸钠组控制了 67% 的 c 波下降 ( $P < 0.05$ )。在脉络膜血流的测量中, 30, 60, 和 120min 的结果显示, TMP 显著增加脉络膜血流。在氧化应激部分, 不同浓度的 TMP 在各种氧化应激损伤中, 对 RPE 都有各种程度的保护作用。

**结论:** 浓度为 1% Tetramethylpyrazine 可以显著保护碘酸钠和氧化应激诱导的 RPE 变性, 增加脉络膜血流, 并可能在 AMD 的治疗中发挥作用。

**关键词:** Tetramethylpyrazine; 碘酸钠; 视网膜色素上皮细胞; 年龄相关性黄斑变性; 脉络膜血流; 氧化应激