

Hyphema due to blunt injury: a review of 118 patients

V Ulagantheran, M S Ahmad Fauzi, S C Reddy

Department of Ophthalmology, Faculty of Medicine,
University of Malaya, Kuala Lumpur, Malaysia

Correspondence to: S C Reddy. Department of Ophthalmology,
UCSI School of Medicine, Bukit Khor, 21600 Marang,
Terengganu, Malaysia. profscreddy@gmail.com

Received:2010-08-23 Accepted:2010-09-09

Abstract

• **AIM:** To determine the causes, associated ocular findings and visual acuity on presentation, complications and visual outcome following treatment in patients of hyphema due to blunt injury

• **METHODS:** A retrospective study was performed in 118 patients with hyphema due to blunt injury admitted in University of Malaya Medical Centre, Kuala Lumpur, Malaysia. The gender, age, race, cause of blunt injury resulting in hyphema, eye involved, vision at admission, other associated ophthalmological findings, complications and visual outcome were noted from the case records of patients. The data were analyzed using SPSS programme.

• **RESULTS:** Males were more predominantly affected (93.2%). Over two-thirds of patients (67.8%) were aged below 30 years. Sports related injury (38.1%) was the most common cause for hyphema. Hyphema disappeared within 5 days in 66.9% of patients. Iris injuries were very commonly associated in the form of mydriasis, sphincter tear and iridodialysis. Associated vitreous haemorrhage was noted in 11.9% of patients. During the hospital stay, secondary haemorrhage was observed in 3.4% of patients. The best corrected vision of 6/18 or better was noted in 85.4% of patients at the last follow-up. The follow-up of these subjects was very poor and thus the incidence of secondary glaucoma could not be established. Moderate blood staining of cornea occurred in 0.8% of patients.

• **CONCLUSION:** Sports related injury is the most common cause of hyphema in Malaysia. Good visual recovery, without serious complications, is possible with appropriate and in-time treatment in hyphema patients due to blunt injury.

• **KEYWORDS:** blunt injury; hyphema; sports injury; blood staining of cornea

DOI:10.3969/j.issn.1672-5123.2010.11.002

Ulagantheran V, Ahmad Fauzi MS, Reddy SC. Hyphema due to blunt injury: a review of 118 patients. *Int J Ophthalmol(Guoji Yanke Zazhi)* 2010;10(11):2049-2053

INTRODUCTION

Injury to the eye is one of the most common causes of unilateral blindness worldwide. Hyphema and concomitant injuries to ocular structures following blunt trauma are not an infrequent cause of presentation to the emergency unit in many hospitals^[1-5]. The causes of poor vision after blunt injury include black ball hyphema, secondary glaucoma, cataract, vitreous haemorrhage, commotio retinae and retinal detachment^[6]. Recognition of factors related to poor visual outcome, appropriate medical therapy, surgical intervention when indicated and careful follow up will help in preserving vision in these patients. Posterior segment injuries^[7], size of hyphema at presentation and retinal damage^[8] have been reported to be significant predictors of poor visual outcome in these patients; while topical steroids and/or cycloplegic medication^[8] and occurrence of secondary haemorrhage^[7,8] did not influence the final visual outcome.

There is no report available in the literature on traumatic hyphema from Malaysia, even though many reports are available on this from different countries in the world^[19]. Therefore, this study was undertaken to determine the cause of blunt injury resulting in hyphema, visual acuity and other ocular findings on presentation, complications and visual outcome in these patients. This information may help in the development of appropriate preventive measures to reduce the morbidity in cases of blunt injuries of eye.

MATERIALS AND METHODS

Subjects This is a retrospective study of 118 patients with traumatic hyphaema due to blunt injury treated in the eye ward of University of Malaya Medical Centre, Kuala Lumpur over a period of ten years. Patients with postoperative hyphema, perforating injury with hyphema, and those who absconded from the ward within a day after admission were not included in this study. After taking history, visual acuity was tested on Snellen chart and the anterior segment was examined with slit-lamp biomicroscope. Intraocular pressure (IOP) was measured with applanation tonometer and fundus examination was done after dilating pupil with 2.5% phenylephrine eye drops using 90 D lens and slit-lamp.

Methods The age, gender, race, type of blunt injury, vision at admission, other ocular findings, grading of hyphema, treatment given, time taken for absorption of hyphema, follow-up period, complications and best corrected vision at last follow-up were noted from the patients' records. Based on the level of blood in the anterior chamber (AC) at

Table 1 Different objects responsible for blunt injury in 118 patients with hyphema

Different objects	n (%)	Different objects	n (%)
Sports related injury		Home injury	
Shuttle cock	25(21.2)	Bottle cap	4(3.4)
Foot ball	5(4.2)	Elastic spring	3(2.5)
Shuttle racquet	4(3.4)	Rubber hosepipe	2(1.7)
Squash ball	4(3.4)	Broom stick	2(1.7)
Airgun plastic bullet	3(2.5)	Wooden stick	2(1.7)
Table tennis ball	2(1.7)	Hit on the door edge	1(0.8)
Golf ball	1(0.8)	Fire cracker	1(0.8)
Tennis ball	1(0.8)	Toothpaste tube	1(0.8)
Industrial/work injury		Assaults injury	
Nail	6(5.1)	Fist/ punch	7(5.9)
Hammer	4(3.4)	Wooden stick	4(3.4)
Wood stick	5(4.2)	Rubber hosepipe	2(1.7)
Grass cutting	3(2.5)	Rubber belt	2(1.7)
Battery explosion	2(1.7)	Motor vehicle accident	
Metal piece	2(1.7)	Fall on the ground	4(3.4)
Bottle cover	2(1.7)	Hit on the door	3(2.5)
Radiator cap	2(1.7)	Fall in the drain	2 (1.7)
Screw driver	2(1.7)		
Wire cable	2(1.7)		

slit-lamp examination, hyphema was graded as microscopic hyphema when no clot was present (circulating red blood cells only), grade I-blood filling < 1/3 of AC, grade II-blood filling < 1/2 of AC, grade III-blood filling more than 1/2 of AC, and grade IV-total hyphema with red or black clots^[1]. The standard protocol of management of hyphema was followed which included admission to the hospital, complete bed rest with restricted ambulation to bath room, elevation of head up to 45 degrees, patching the affected eye with rigid shield, using topical cycloplegics (homatropine 20g/L eye drops *bd*), corticosteroid (dexamethasone 1g/L eye drops *qid*), and Tab. Vitamin C 500mg daily. Eyes with raised intraocular pressure were treated with timolol 5g/L eye drops *bd*, and Tab. Acetazolamide 250mg *tds*. When the IOP was high (more than 30mmHg) inspite of glaucoma medications, IV mannitol was given. Surgical evacuation of the blood was performed in eyes with total hyphema and uncontrolled intraocular pressure with medication. Patients were monitored of blood level in the anterior chamber, visual acuity and intraocular pressure daily during their stay in the hospital. Secondary haemorrhage was diagnosed if the size of hyphema was increased or if fresh blood was present in the anterior chamber.

Gonioscopy was performed during the examination in first follow-up visit. The angle appearance was noted according to Shaffer classification from grade 0 to grade IV angle and abnormalities such as angle recession and peripheral anterior synechiae were also noted. In patients where gonioscopy was not done, it was listed as not done while noting down the findings from records. The above data were tabulated and

analyzed using SPSS programme.

RESULTS

Out of 118 patients, majority were males (110 cases, 93.2%). Age: 11-20, 21-30, 31-40, 41-50, 51-60, 61-70 and 71-80 years were 29 cases (24.6%), 51 cases (43.2%), 20 cases (17.0%), 10 cases (8.5%), 6 cases (5.1%), 1 case (0.8%) and 1 case (0.8%), respectively. Over two-thirds of patients (67.8%) were young and below 30 years of age. The mean age of the patients was 21.6 years (range 12-78 years). 39.5% (n=46) were Malays, 40 cases (33.5%) were Chinese; 16 cases (13.6%) were Indonesian; 11 cases (9.3%) were Indian and 5 cases (3.8%) were Bangladesh. Hyphema was present in one eye only in all the patients in our study, and 63 eyes (53.3%) were in the right; 55 eyes (46.7%) were in the left. Sports injuries (n=45, 38.1%) were the most common cause of hyphema, the others being industrial (n=30, 25.4%), home (n=16, 13.5%), assault (n=15, 12.7%) and motor vehicle accidents (n=9, 7.6%). The details of different objects responsible for blunt injury and resulted in hyphema are shown in Table 1. The visual acuity at presentation ranged from perception of light to 6/6; 29 (24.5%) of them had poor vision (<6/60); 6 cases (5.0%) had perception of light (PL), 9 cases (7.6%) had hand movements (HM), 6 cases (5.0%) had counting fingers (CF), 8 cases (6.7%) had 3/60, 10 cases (8.4%) had 6/60, 5 cases (4.2%) had 6/36, 8 cases (6.7%) had 6/24, 10 cases (8.4%) had 6/18, 17 cases (14.3%) had 6/12 and 14 cases (11.8%) had 6/9. Vision was normal in 25 (21.0%) patients. The IOP at admission ranged between 8 and 42mmHg. The IOP of 20mmHg or below was seen in 91(77.1%)

Table 2 Associated ocular findings in 118 patients with hyphema

Ocular findings *	n(%)	Ocular findings *	n(%)
Eyelids		Iris and pupil	
Oedema	43(36.4)	Mydriasis	65(55.1)
Echymosis/haematoma	38(32.2)	Sphincteric tear	59(50.0)
Laceration	2(1.7)	Iridodialysis	11(9.3)
Conjunctiva		Miosis	2(1.7)
Chemosis	50(42.4)	RAPD	2(1.7)
Sub conj. haemorrhage	16(13.6)	Lens	
Cornea		Cataract	12(10.2)
Oedema	52(44.1)	Phacodonesis	4(3.4)
Abrasion	42(35.6)	Dislocation	3(2.5)
Partial laceration	2(1.6)	Retina	
		Comotio retinae	37(31.0)
		Vitreous haemorrhage	14(11.9)
		Macular striae	2(1.7)
		Macular hole	1(0.8)
		Choroidal detachment	1(0.8)

* Some patients had more than one finding; RAPD: Relative afferent pupillary defect

patients; 19 (16.1%) had 21-30mmHg and 8 (6.7%) had 31mmHg and above (secondary glaucoma in 27 (22.8%) patients). In addition to hyphema, there were associated findings in the globe and adnexa (Table 2). Echymosis/haematoma was seen in 32.2% of patients while subconjunctival haemorrhage was seen in 13.6% and bleeding into the vitreous in 11.9% of patients. More than one sign were present in some of the patients. Hence the total number of patients are much more in this table. There was no fracture of orbital bones in any of the patients. The most common form of hyphema in our study was grade I, seen in 66(55.9%) patients, Grade II in 20(16.9%), Grade III in 12(10.2%), Grade IV in 10(8.5%), and microscopic in 9(7.6%). Patients with grade III/IV hyphema had associated periorbital haematoma in 14 patients, iridodialysis in 3 patients, and traumatic mydriasis in 18 patients. Secondary haemorrhage was seen on second day after admission in 2(1.6%) patients and on third day in 1(0.8%) patient, which resolved completely with bed rest and treatment. There were sphincter tears in these patients and all were shuttle cock injuries. All patients were treated as per the standard protocol as mentioned in the methods. Surgical intervention was needed only in 2 elderly patients of grade IV hyphema (vision only hand movements) in whom the IOP did not come down to normal level in spite of medical treatment. Paracentesis and evacuation of blood clot was done in both patients. The IOP was controlled with medical treatment. The best corrected vision improved to 6/18 in one patient. However, the other patient developed moderate blood staining of cornea and vision improved to only 6/60. This patient had cataract in the affected eye; following extracapsular cataract extraction and posterior chamber intraocular lens implantation, the best

corrected vision improved to 6/24 only. The IOP on discharge was 20mmHg or below in 115(97.5%) patients and between 21 and 30mmHg in 3(2.5%) patients. The IOP came to normal reading with continued glaucoma medication in these 3 patients during the follow up examination after 3 weeks. The time taken for absorption of hyphema ranged between 2 and 20 days; in two-thirds of patients (79, 66.9%) hyphema was absorbed within 5 days. Five out of 22 patients with grade III/IV hyphema stayed for 15 days or more in the hospital. All these patients were discharged from the hospital only after complete absorption of hyphema. Gonioscopy findings were available in 69 patients only; the high default rate of follow up examinations was the probable cause for this. The angle of the anterior chamber was normal in 50 out of 69 (72.4%) patients; angle recession was 90° in 15 (21.7%); 180° in 3 (4.3%); and 360° in 1 (1.4%) patients. All the patients with angle recession sustained shuttle cock injuries. Follow-up of patients was very poor; ranging from 1 to 56 months. The follow up was for 1 month in 84 (71.2%) patients, for 6 months in 70 (59.3%) and for 12 months in 41 (34.7%) patients. Secondary glaucoma was noted in 8 out of 19 patients with angle recession. Since the gonioscopy was not done in all the patients in the present study and long term follow up was very poor, this figure can not be taken into account regarding then incidence of secondary glaucoma. The best corrected vision recorded at the last follow up ranged between counting fingers (n = 3, 2.5%) and 6/6 (n = 66, 55.9%); 15 (12.7%) had 6/9; 12 (10.1%) had 6/12; 8 (6.7%) had 6/18; 3 (2.5%) had 6/24; 3 (2.5%) had 6/36; 4 (3.4%) had 6/60; and 4 (3.4%) had 4/60; and 7 (5.9%) of them had poor vision (<6/60) and 101(85.6%) patients had vision of 6/18 or better.

DISCUSSION

Kuala Lumpur is the capital city of Malaysia. There are three tertiary government hospitals (two university hospitals and one ministry of health general hospital) and one private eye hospital where eye emergencies are treated. In addition, there are many private eye specialists practicing in this city. The probable reason for small number of patients in this study over a period of ten years could be the availability of many eye care facilities in the city. The higher frequency of hyphema in males in our study is because of more males working in different industries, their participation in sports and their involvement in assaults/accidents. The male to female ratio in our study (2.5:1) is similar to that reported from Nigeria^[5,6]. However, much higher frequency of this ratio (8:1) was reported from south Korea^[7]. Two-thirds (67.8%) of patients in our study were young (below 30 years) and a similar figure was reported from UK (77%)^[3]. However, similar age patients were reported to be in much higher percentage of patients from Nigeria (90%^[6] and 96%^[5]). In a prospective study of 5671 cases of eye injuries presented to the eye casualty, Macewen^[7] from UK reported hyphema in 31(0.6%) cases. In their study, the eye injuries resulting in hyphema occurred during different activities such as sports (21 cases), domestic (5), at work (3), assault (10) and unknown cause (1 case).

In our study, sports/ games related injuries were the most common cause for hyphema (38.1%), of which shuttle badminton game accounted for 64.4% (29 out of 45 cases). The ocular injury in this game occurs commonly with round ended shuttle cock in the smash hit by opponent player in the single game, while the injury is due to long stick end of the racket of the partner player or with shuttle cock hit by the opponent player in the doubles game. Getting injury to the eye is an unacceptable price while playing this game. Wearing the protective glasses during play like in squash game can prevent ocular injuries. Sports/play related injuries reported from UK (49.2%)^[3], Nigeria (42.1%)^[5] and Pakistan (66.6%)^[4] are much higher than our study (38.1%). However, a much lower figure (9.1%)^[6] has been reported in a recent study of 472 cases of hyphema from Nigeria. Industrial accidents/accidents at work place responsible for hyphema were more common in foreigners (Indonesians, Bangladeshis) because majority of the construction workers/people working in factories are from Indonesia and Bangladesh. Kearns^[3] reported accident at work was the cause of hyphema in 9.9% of patients in UK, while a less frequency of the same (7.6%)^[6] was observed in Nigeria. Domestic causes/injury at home was responsible for hyphema in 13.5% of the cases in our study, while the same was higher (17.8%)^[3] in UK patients. However, a much lower figures (7.9%^[5] and 2.5%^[6]) were reported in Nigerian patients. Shaking the carbonated

drinks bottle before opening the cap to release the gas faster, is the most common cause of bottle cap eye injuries at home. Bottle cap injuries were noted in 3.4% of patients in our study while the same was reported in 2.7% of patients in Seoul by Cho *et al*^[7]. Assaults injury with different objects was responsible for hyphema in 12.7% of patients in our study while the same was reported in 20.7% of patients in UK by Kearns^[3]. Motor vehicle accidents resulted in hyphema in 7.6% of cases in our study, while the traffic accidents as the cause of hyphema has been reported to be in 1.6% of cases by Cho *et al*^[7]. Secondary haemorrhage was seen on second day after trauma in 3.4% of patients, while the same was reported to be varying between 3.8%^[7], 4.1%^[3], 6.7%^[2], 8.7%^[10] and 9.2%^[5] of patients in different countries. Ng *et al*^[10] demonstrated that the size of hyphema on presentation and the presence of retinal damage did not affect the probability of secondary haemorrhage. The incidence of secondary haemorrhage was found to decrease significantly with the use of topical steroid, but did not appear to be influenced by the use of cycloplegics. This emphasizes the therapeutic importance of topical steroid in the treatment of hyphema due to blunt trauma.

The main treatment is bed rest and topical medical therapy to prevent iritis. Visual acuity and intraocular pressure are to be monitored daily during the hospitalization of the patient. Surgical evacuation of the blood has to be done when the IOP is 50 mm Hg for >5 days, or 30 mm Hg for >7 days inspite of maximal anti glaucoma medications; when there is no sign of absorption of blood within 3-4 days after injury in patients with total hyphema and when there is impending blood staining of cornea. The optimum duration of surgery is 5th to 7th day after injury for the following reasons; (1) if the IOP is normal, blood staining of cornea is unlikely to occur after 5th day of trauma; (2) too early attempts to remove the blood clot may in fact cause further bleeding because of manipulations in the anterior chamber, sudden lowering of IOP and dislodgement of clot; (3) surgery after 7 days of injury would be difficult due to firm adhesions and fibrosis; (4) new blood vessels and organization of clots occur within the first 5 days after trauma; (5) secondary haemorrhage mostly occurs in the first 4 days after trauma^[11]. The best corrected vision of 6/18 or better in 85.6% of patients in our study indicates the beneficial effect of appropriate and in-time treatment in patients of hyphema due to blunt trauma. The causes of poor vision in the rest of patients were the effects of associated retinal findings such as commotion retinae, macular edema and scarring, macular hole and secondary glaucoma. Wearing protective glasses during shuttle badminton, squash games and at work places, supervision of children while playing, dissemination of preventive methods of eye injuries through media and in the schools/colleges will help to reduce the ocular morbidity in young patients.

REFERENCES

- 1 Edwards WC, Layden WE. Traumatic hyphema: A report of 184 consecutive cases. *Am J Ophthalmol* 1973; 75(1): 110-116
- 2 Yospaiboon Y, Sangveejit J, Suwanwatana C. Traumatic hyphema: clinical study of 149 cases. *J Med Assoc Thai* 1989;72(9): 520-526
- 3 Kearns P. Traumatic hyphema: a retrospective study of 314 cases. *Br J Ophthalmol* 1991; 75(3): 137-141
- 4 Jan S, Khan S, Mohammad S. Hyphema due to blunt trauma. *J Coll Physicians Surg Pak* 2003; 13(7): 398-401
- 5 Amoni SS. Traumatic hyphema in Kaduna, Nigeria. *Br J Ophthalmol* 1981; 65(6): 439-444
- 6 Ashaye AO. Traumatic hyphema: a report of 472 cases. *BMC Ophthalmol* 2008;8:24
- 7 Cho J, Jun BK, Lee YJ, Uhm KB. Factors associated with the poor final visual outcome after traumatic hyphema. *Korean J Ophthalmol* 1998; 12(2): 122-129
- 8 Ng CS, Sparrow JM, Strong NP, Rosenthal AR. Factors related to the final visual outcome of 425 patients with traumatic hyphema. *Eye (London)* 1992; 6(Pt 3): 305-307
- 9 Macewen CJ. Eye injuries: a prospective study of 5671 cases. *Br J Ophthalmol* 1989;73(11): 888-894
- 10 Ng CS, Strong NP, Sparrow JM, Rosenthal AR. Factors related to the incidence of secondary haemorrhage in 462 patients with traumatic hyphema. *Eye (London)* 1992; 6(Pt 3): 308-312
- 11 Low RC. The management of traumatic hyphema. *Singapore Med J* 1975;16(2):122-127

钝伤性前房出血 118 例回顾分析

V Ulagantheran, M S Ahmad Fauzi, S C Reddy
(作者单位:马来西亚吉隆坡,马来亚大学医学院眼科)
通讯作者: S C Reddy. profscreddy@gmail.com

摘要

目的:寻找因钝器损伤导致前房积血的原因、视力损伤等相关情况,并探讨治疗后的并发症及预后视力情况。

方法:回顾性分析在马来亚大学医疗中心(马来西亚吉隆坡)治疗的因钝器损伤导致前房积血 118 例患者。从患者的病情记录获取患者的性别、年龄、种族、钝伤导致的积血原因、眼附属物、入院视力、相关眼科情况、并发症和视力结果等资料,并采用 SPSS 软件进行分析。

结果:男性受累居多,占 93.2%,有 2/3 患者(67.8%)年龄 < 30 岁。运动相关的伤害(38.1%)是最常见的前房积血原因。66.9% 的患者 5d 内前房出血消失。虹膜损伤非常普遍,包括括约肌撕裂,虹膜根部脱离等,而并发玻璃体出血患者约占 11.9%。住院期间,二次观察出血的患者为 3.4%。在最后的随访中视力 $\geq 6/18$ 的最佳矫正视力占 85.4%。这些患者随访不到位导致继发性青光眼的发病率无法确定。中度角膜血染患者约占 0.8%。

结论:在马来西亚,运动相关的损伤是前房出血最常见的原因。适当和及时的治疗能使钝伤性前房积血患者具有良好的视力,且无严重并发症发生。

关键词:钝伤;前房出血;运动损伤;角膜血染

Great News: IJO was accepted for coverage in SCIE

International Journal of Ophthalmology——IJO(English edition), established in 2008, was applied for the indexation of SCI in 2009, and on September 21, 2010 we received an email from an editor of SCI, informing us that our journal was accepted for the coverage in SCIE. This is the only ophthalmological periodical in China indexed by the most authoritative international searching organization——SCIE.

Hereby, we would like to give all the ophthalmologists and authors both from home and abroad our special thanks for the valuable guidance and great support for our journal, and we sincerely hope you success in your career, and let us make a great achievement together!

Contributions and guidance are greatly welcomed!

Editorial Office of IJO
2010-10-21