

# Correlation of serum insulin like growth factor-1 with retinopathy in Malaysian pregnant diabetics

S Adzura<sup>1</sup>, M Muhaya<sup>2</sup>, M Normalina<sup>3</sup>, A M Zaleha<sup>4</sup>, W P Sharifa Ezat<sup>5</sup>, I Tajunisah<sup>6</sup>

<sup>1</sup>Department of Ophthalmology, Kulliyyah of Medicine, International Islamic University, Kuantan, Pahang, Malaysia

<sup>2</sup>Department of Ophthalmology, Faculty of Medicine, National University of Malaysia, Kuala Lumpur, Malaysia

<sup>3</sup>Department of Ophthalmology, Faculty of Medicine, MARA University of Technology, Kuala Lumpur, Malaysia

<sup>4</sup>Department of Obstetrics & Gynaecology, Faculty of Medicine, National University of Malaysia, Kuala Lumpur, Malaysia

<sup>5</sup>Department of Community Health, Faculty of Medicine, National University of Malaysia, Kuala Lumpur, Malaysia

<sup>6</sup>Department of Ophthalmology, Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia

**Correspondence to:** Tajunisah Iqbal. Department of Ophthalmology, Faculty of Medicine, University of Malaya, 50603 Kuala Lumpur, Malaysia. tajun69@yahoo.com

Received:2010-10-21 Accepted:2011-03-14

## Abstract

• **AIM:** To study the association of serum insulin-like growth factor-1 (IGF-1) with diabetic retinopathy.

• **METHODS:** Serum IGF-1 levels were measured in 25 pregnant diabetic patients and 25 pregnant non-diabetic patients who were matched for age, ethnicity, parity and period of gestation. Fundus examination was performed in both groups at 28, 32 and 36 weeks of gestation.

• **RESULTS:** The serum IGF-1 level was significantly elevated in pregnant diabetics compared to pregnant non-diabetics [ $366 \pm 199 \mu\text{g/L}$  vs  $184 \pm 89 \mu\text{g/L}$  ( $P = 0.0001$ ) at 24 weeks,  $535 \pm 251 \mu\text{g/L}$  vs  $356 \pm 89 \mu\text{g/L}$  ( $P = 0.007$ ) at 32 weeks and  $404 \pm 166 \mu\text{g/L}$  vs  $264 \pm 113 \mu\text{g/L}$  ( $P = 0.003$ ) at 36 weeks of gestation]. The pregnant diabetics with established diabetes had significantly higher IGF-1 level than gestational diabetes at 28, 32 and 36 weeks of gestation. The serum IGF-1 level in pregnant diabetics with retinopathy was significantly higher than in those without retinopathy at all periods of gestation.

• **CONCLUSION:** Increased serum IGF-1 in pregnancy may increase the risks for retinopathy.

• **KEYWORDS:** diabetic retinopathy; pregnancy; serum IGF-1 level

DOI:10.3969/j.issn.1672-5123.2011.04.003

Adzura S, Muhaya M, Normalina M, Zaleha AM, Sharifa Ezat WP, Tajunisah I. Correlation of serum insulin like growth factor-1 with retinopathy in Malaysian pregnant diabetics. *Guoji Yanke Zazhi (Int J Ophthalmol)* 2011;11(4):573-576

## INTRODUCTION

In South East Asia, the prevalence of blindness has been reported to be 1.2% in Indonesia, 1.1% in Thailand and 0.8% in Vietnam<sup>[1]</sup>. The prevalence of blindness among the Malaysian adult between the age of 30 and 70 was 0.29%, with 14.6% due to diabetic retinopathy<sup>[2]</sup>. In pregnant diabetic women, the prevalence of diabetic retinopathy is high, ranging from 10% to 27%. Based on the Diabetic Control and Complication Trial Research Group (DCCT) results, pregnant diabetic women have a 1.63-fold greater risk of any worsening of retinopathy from before pregnancy to during pregnancy in the intensive treatment group, the risk was 2.48-fold greater for pregnant vs non pregnant women in the conventional group<sup>[3]</sup>. The retinopathy progresses markedly among the IDDM and NIDDM compared with gestational diabetes. Diabetic retinopathy is known to progress during pregnancy but the exact mechanism is still unknown. Clinical evidence has suggested that insulin-like growth factor (IGF) system may be involved in the development of retinopathy in pregnancy. Elevated vitreous and aqueous IGF-1 levels are strongly linked to proliferative diabetic retinopathy and other neovascular disorders secondary to retinal ischemia<sup>[4]</sup>. Lauszus *et al*<sup>[5]</sup> showed that despite good glycemic control in pregnancy, an increase in circulating IGF-1, takes place concomitantly with progression of diabetic retinopathy. This is possible because serum and vitreous IGF-1 concentration have been shown to correlate, probably due to leakage of IGF-1 from the blood stream. This study was undertaken to compare the level of serum IGF-1 between pregnant diabetic patients (DP) and pregnant non-diabetic patients (NDP) at different periods of gestation during the third trimester. In this study, we also measured the level of serum IGF-1 in pregnant women with established diabetes (NIDDM and IDDM) compared to those with gestational diabetes. Since the level of serum IGF-1 has been shown to be high in pregnant women with diabetic retinopathy, we also studied the significance between diabetic retinopathy and serum IGF-1 level in pregnant diabetics.

## MATERIALS AND METHODS

**Materials** This case-controlled, prospective study included patients with established diabetes, gestational diabetes and non-diabetics attending the Obstetrics and Gynaecology Clinic, National University of Malaysia from Jan. 2004 to Jan. 2005. Patients meeting the inclusion criteria were included into this study and they were selected based on method of universal sampling. Pregnant women with established diabetes and/or gestational diabetes mellitus were

included in the study. Established diabetes is defined in a woman who becomes pregnant and who is known to have diabetes mellitus (IDDM or NIDDM), which antedates the current pregnancy. Gestational diabetes mellitus is a carbohydrate intolerance resulting in hyperglycemia of variable severity with onset of first recognition during pregnancy. It is diagnosed by oral glucose tolerance test (OGTT) within gestational age of  $28 \pm 2$  weeks and  $36 \pm 2$  weeks. Any pregnant woman who met the WHO criteria for diabetes mellitus or impaired glucose tolerance was classified as having gestational diabetes mellitus. Normal, non-diabetic pregnant women (NDP), age, ethnicity and parity matched at third trimester within gestational age of  $28 \pm 2$  weeks and  $38 \pm 2$  weeks were taken as the control group. Any patients with previous history of laser treatment; with concurrent other endocrine problems such as hyperthyroidism and hypertension; with ocular pathology such as glaucoma, high myopia (more than -6D) and cataracts; with end stage renal failure and nephrotic syndrome were excluded. Informed consent was obtained from all patients upon ethical committee clearance, and the study was conducted in accordance with the Declaration of Helsinki and subsequent revisions thereof. Based on these two criteria, a total of 50 patients were examined, which included 25 pregnant diabetics (DP) (established diabetes and gestational diabetes) and 25 pregnant non diabetic patients (NDP) who were matched according to age, ethnicity, gestation and parity. There was no significant difference in mean age ( $P = 0.4$ ), ethnic distribution ( $P = 0.79$ ) and parity status ( $P = 0.24$ ) between the DP and NDP groups. There were 21 (42%) Malays, 21 (42%) Chinese, 5 (10%) Indians and 3 (6%) patients from other races. The mean age of patients was  $31.56 \pm 6.32$  years in DP group and  $29.44 \pm 3.94$  years in NDP group. Out of the 25 patients in DP group, 17 (34%) were established diabetics (Type 2 diabetes,  $n = 15$  and Type 1 diabetes,  $n = 2$ ), and 8 (16%) patients were gestational diabetics. Seven patients were primigravida (14%) and 18 (36%) were multigravida. In NDP group, 9 (18%) patients were primigravida and 16 (32%) patients were multigravida. In DP group, the duration of diabetes mellitus was between 1 to 18 years with a median of one year. Among the established diabetics group, there were 2 out of 17 patients detected to have diabetic retinopathy at first fundus examination at 28 weeks of gestation and both of them were IDDM. The first patient had asymmetrical diabetic retinopathy changes where she developed proliferative retinopathy in the left eye and mild non-proliferative changes in the right eye. She had 10 years history of IDDM (since age of 16 years), whereas the diabetes was poorly controlled. Pan retinal photocoagulation (PRP) treatment for the left eye was given over four sessions with a total number of 2569 burns. She unfortunately developed two episodes of vitreous haemorrhages in the left eye during the third trimester despite laser PRP. In view of the poorly controlled diabetes, an emergency lower segment caesarean section (LSCS) was performed at 36 weeks gestation. She gave birth to a healthy baby boy with a birth weight of 3.4kg. One month after delivery, visual acuity of

the left eye improved to 6/9 and fundus examination showed a left quiescent proliferative diabetic retinopathy. The second patient with retinopathy in the established diabetes group had mild non-proliferative changes in both eyes. She had 8 years history of IDDM (since age of 16 years). In contrast to the first patient, her diabetes was well controlled during her pregnancy. She gave birth to a healthy baby girl with a birth weight of 3.6kg via spontaneous vaginal delivery. Two months after delivery, the fundus examination showed no progression of her retinopathy bilaterally.

**Methods** All patients who fulfilled the inclusion criteria underwent examination at 28 weeks of gestation that consisted of height, weight and blood pressure measurement, assessment of best-corrected vision acuity, followed by slit-lamp biomicroscopic examination and intraocular pressure reading. The pupil was dilated with tropicamide 10g/L and phenylephrine 25g/L eye drops. Fundus was examined by indirect ophthalmoscopy and posterior pole examination was performed with 90 diopter lens when necessary. Fundus photography was performed with Topcon Retinal Camera TRC-50IX with Image Net Digital Imaging System. The retinopathy was graded and findings were entered in the data collection sheet. The fundus examination was repeated at 32 weeks and 36 weeks of gestation. The patients' blood samples were taken using 21G needle; 3mL of blood was syringed out from each patient for measurement of IGF-1 during each visit. The blood was centrifuged to separate the serum from other components using the centrifuge machine at the rate of 3000r/min for 10 minute duration. The sample was then kept in the freezer at  $-20^{\circ}\text{C}$  until adequate sample amount was available for analysis. IGF-1 was measured using Automated Chemiluminescent Immunoassays on the IMMULITE<sup>®</sup> machine analyzer<sup>[6]</sup>. This technique was compatible with Nichols Advantage<sup>™</sup> assay for serum IGF-1 level measurement<sup>[7]</sup>.

**Statistical Analysis** The difference in mean of serum IGF-1 between DP group and NDP group and between pregnant women with diabetic retinopathy and pregnant women with no diabetic retinopathy was tested using Mann-Whitney *U* test. The difference in mean of serum IGF-1 level between established diabetes and gestational diabetes groups was tested using Kruskal-Wallis test. The association between diabetic retinopathy and IGF-1 level was tested using Chi-square with Fisher's exact test.  $P < 0.05$  was considered statistically significant.

## RESULTS

**Serum IGF-1 in DP and NDP** There was statistically significant difference in serum IGF-1 level between DP group compared to NDP group ( $366 \pm 199\mu\text{g/L}$  vs  $184 \pm 89\mu\text{g/L}$ ,  $P = 0.0001$ ) at 24 weeks, ( $535 \pm 251\mu\text{g/L}$  vs  $356 \pm 89\mu\text{g/L}$ ,  $P = 0.007$ ) at 32 weeks and ( $404 \pm 166\mu\text{g/L}$  vs  $264 \pm 113\mu\text{g/L}$ ,  $P = 0.003$ ) at 36 weeks of gestation. There was significant increase in serum IGF-1 level from 28 to 32 weeks followed by significant decline of serum IGF-1 level from 32 to 36 weeks of gestation (0.0001 in both groups).

**Serum IGF-1 between established diabetes and gestational diabetes** The diabetic pregnant (DP) group was further

divided into established diabetes ( $n = 17$ ) and gestational diabetes mellitus ( $n = 8$ ). There was a statistically significant difference in serum IGF-1 level between established diabetes and gestational diabetes ( $392 \pm 217 \mu\text{g/L}$  vs  $311 \pm 153 \mu\text{g/L}$ ,  $P = 0.01$ ) at 28 weeks, ( $558 \pm 278 \mu\text{g/L}$  vs  $498 \pm 175 \mu\text{g/L}$ ,  $P = 0.025$ ) at 32 weeks and ( $416 \pm 184 \mu\text{g/L}$  vs  $377 \pm 130 \mu\text{g/L}$ ,  $P = 0.012$ ) at 36 weeks of gestation.

**Serum IGF-1 in DP with vs without diabetic retinopathy**

There was statically significant difference in serum IGF-1 level between pregnant diabetic patients with diabetic retinopathy compared to pregnant diabetic patients with no diabetic retinopathy ( $541 \pm 267 \mu\text{g/L}$  vs  $264 \pm 169 \mu\text{g/L}$ ,  $P = 0.03$ ) at 28 weeks, ( $937.5 \pm 112 \mu\text{g/L}$  vs  $425 \pm 184 \mu\text{g/L}$ ,  $P = 0.000$ ) at 32 weeks and ( $645 \pm 282 \mu\text{g/L}$  vs  $322 \pm 141 \mu\text{g/L}$ ,  $P = 0.004$ ) at 36 weeks of gestation. However, There was no positive association between diabetic retinopathy and serum IGF-1 level at different periods of gestation (at 28 weeks ( $P = 0.49$ ), 32 weeks ( $P = 0.225$ ) and 36 weeks of gestation ( $P = 0.49$ )) (Table 1).

**DISCUSSION**

The major finding in this study was a significantly higher difference in serum IGF-1 level in pregnant diabetic group compared to pregnant non-diabetic group during the third trimester of pregnancy. Serum IGF-1 in pregnant diabetic group was doubled compared to pregnant non-diabetic group at 28 weeks of gestation, with 1.5 fold increases at 32 and 36 weeks of gestation. Interestingly, the pattern of serum IGF-1 level showed a significant rise from 28 weeks to 32 weeks gestation in both groups before showing a significant decline from 32 weeks to 36 weeks gestation in both groups. The pattern of serum IGF-1 rise observed in this study is similar to another study by Hayati *et al* [8] who documented a significantly higher concentration of serum IGF-1 in diabetic pregnancy compared to controls, both at 28 and 36 weeks of gestation. Lauszus *et al* [5] also reported similar serum IGF-1 pattern changes, where the level was increased with increased gestational age until a plateau was reached at 34 weeks of gestation followed by a decrease from weeks 36 to 38 and full normalization after delivery.

The pathophysiology for deranged serum IGF-1 in pregnant diabetics is explained by the role of human placental growth hormone (hPGH) which gradually replaces the pituitary growth hormones from week 20 onward. Production of maternal IGF-1 is regulated by placental hormone via hypothalamic-growth hormone-IGF-1 axis. The pituitary growth hormone regulates the secretion of IGF-1, which, in turn, exerts negative feedback action on growth hormone at hypothalamic pituitary level. During the latter half of pregnancy, the growth hormone-IGF-1 axis is inhibited by large amount of oestrogen. The larger increase in placental growth hormone exerts an inhibitory effect on growth hormone secretion mediated by placental growth hormone on the hypothalamus and pituitary. The hPGH is secreted by placenta into the maternal circulation and stimulates maternal IGF-1 production in the liver. IGF-1 level do not vary significantly during the first weeks of gestation, but then increase

**Table 1 Outcome cross tabulation of serum IGF-1 pregnant diabetic women**  $n = 25$

Gestation period (Median, g/L)	Diabetic retinopathy		
	Yes	No	
28 weeks	<224	0	25
	>224	2	23
32 weeks	<404	0	25
	>404	2	23
36 weeks	<336	0	25
	>336	2	23

gradually from  $165 \pm 445 \text{mg/L}$  at about 24-25 weeks of gestation, and reach levels of  $330.5 \pm 63.5 \text{mg/L}$  at 32 weeks of gestation. Increase in maternal serum IGF-1 will result in development of insulin resistance during pregnancy followed by hyperglycemia towards the second and third trimester. In diabetic pregnancy, there is further increase in the production of maternal serum IGF-1 through stimulation of IGF-1 mRNA from the term placenta. This excessive production of maternal serum IGF-1 production is part of adaptive response of the adverse environmental such as hyperglycemia in pregnancy complicated with diabetes.

Based on the above observation, measuring serum IGF-1 in third trimester may help us to identify pregnant women who are at a risk of developing diabetes in pregnancy especially those who belong to high risk groups such as over weight, strong family history of diabetes, previous history conceiving and delivering a big baby and previous history of congenital malformation.

Gestational diabetes is defined as glucose intolerance that begins or first detected during pregnancy. The WHO oral glucose tolerance test (OGTT) criteria is the most commonly used to diagnose this condition. Debates continue as to whether gestational diabetes represents a pregnancy-induced state of glucose intolerance or a state of pre existing insulin resistance that would become evident in the future regardless of pregnancies. There are many similarities between gestational diabetes and the syndrome of insulin resistance in non pregnant subjects [9]. Since this study had shown that IGF-1 was higher in established diabetes group compared to gestational diabetes, it could be proposed that both the combination of serum IGF-1 and the WHO OGTT criteria could be used to differentiate true gestational diabetes from chronic diabetics who become pregnant whom are mistakenly labelled as gestational diabetics. Thus, proper screening prior to pregnancy or while planning for pregnancy is important so as not to miss the group of people who are asymptomatic prior to pregnancy. This study also showed that the serum IGF-1 level in pregnant women with diabetic retinopathy was significantly higher compared to those with no diabetic retinopathy. Lauszus *et al* [5] and Rozita *et al* also supported this finding where they found a significant rise in serum IGF-1 among diabetic patients with diabetic retinopathy compared with diabetic patients without diabetic retinopathy. It has been hypothesized that a rise in serum IGF-1 level may be related to

vascular microangiopathy and risk of end organ damage development such as diabetic retinopathy and diabetic nephropathy. The pathophysiology behind this microangiopathy was observed in a study by Grant *et al* who reported a high IGF-1 level in the vitreous body of diabetic patients with diabetic retinopathy compared to non diabetic patients. That study also suggested that serum and vitreous IGF-1 concentration were related probably due to leakage of IGF-1 from blood stream. Cummings *et al*<sup>[10]</sup> reported a strong association between urinary IGF-1 and microalbuminuria in patients with type 1 diabetes. The same study also found a significant correlation between plasma IGF-1 and urinary IGF-1 in type 1 diabetes patients. Based on this, one could postulate that in clinical setting, measurement of serum IGF-1 may be able to predict those pregnant diabetics who are at high risk of developing diabetic retinopathy and other end organ damage due to diabetic microangiopathy. In the future, it is hoped that IGF-1 may serve, like VEGF, as a potential therapeutic target for the treatment of intraocular neovascularization especially in diabetic retinopathy. However further studies and clinical trials are required to achieve this. This study has shown that the mean serum IGF-1 level is elevated in diabetic pregnancy compared to non diabetic pregnancy during the third trimester and that the level was markedly higher in pregnant diabetics with diabetic retinopathy. Thus it is recommended that women who are planning for pregnancy should undergo proper diabetic screening since increased levels of IGF-1 in the pregnant state may increase the risks for diabetic retinopathy which is a major visual morbidity.

#### REFERENCES

- 1 Rema M, Mohan V. Retinopathy at diagnosis among young Asian diabetic patients; ASDIAB Study Group. *Diabetes* 2002;51(Suppl 2):A206-207
- 2 Zainal M, Ismail SM, Ropilah AR, Elias H, Arumugam G, Alias D, Fathilah J, Lim TO, Ding LM, Goh PP. Prevalence of blindness and low vision in Malaysian population; results from the National Eye Survey 1996. *Br J Ophthalmol* 2002;86(9):951-956
- 3 Diabetes Control and Complications Trial Research Group. Effect of pregnancy on microvascular complications in diabetes control and complication trial. *Diabetes Care* 2000;23:1084-1091
- 4 Burgos R, Carlos M, Canton A, Ana C, Cristina H, Jorge M, Rafael S. Vitreous level of Insulin like growth factor-IGF-I, insulin like growth factor-binding protein-IGF-I and insulin like growth factor-binding protein-IGF-3 in proliferative diabetic retinopathy. *Diabetes Care* 2000;23:80-83
- 5 Lauszus FF, Klebe JG, Bek T, Flybjerg A. Increase serum IGF-I during pregnancy is associated with progression of diabetic retinopathy. *Diabetes* 2003;52(3):852-856

- 6 Wen J, Bostanian Z, Lei JD, Whitters E, Sustarsic D, Unver E. Automated chemiluminescent immunoassays of insulin-like growth factor I and insulin-like growth factor binding protein 3 on the IMMULITE analyzer. *Clin Chem* 2002;48(Suppl S6):A128
- 7 Valsamis J, Vankrieken J. Comparison of the IMMULITE 2000 AFP assay to three other assay for alphafetoprotein on amniotic fluid samples. Poster 250. An association for Clinical Chemistry 52<sup>nd</sup> Annual meeting and Clinical Lab Exposition 2000
- 8 Hayati AR, Cheah FC, Yong JF, Tan AE, Norizah WM. The role of serum insulin-like growth factor I (IGF-I) in neonatal outcome. *J Clin Pathol* 2004;57(12):1299-1301
- 9 White P. Diabetes mellitus in pregnancy. *Clin Perinatol* 1974;1:331-347
- 10 Cumming EA, Sochett EB, Dekker MG, Lawson ML, Daneman D. Contribution of growth hormone and IGF-I to early diabetic nephropathy in type 1 diabetes. *Diabetes* 1998;47(8):1341-1346

## 糖尿病孕妇血清胰岛素样生长因子-1 与视网膜病变的关系

S Adzura<sup>1</sup>, M Muhaya<sup>2</sup>, M Normalina<sup>3</sup>, A M Zaleha<sup>4</sup>, W P Sharifa Ezat<sup>5</sup>, I Tajunisah<sup>6</sup>

(作者单位:<sup>1</sup>马来西亚彭亨关丹,国际伊斯兰大学 Kulliyyah 医学院眼科;<sup>2</sup>马来西亚吉隆坡,马来西亚国立大学医学院眼科;<sup>3</sup>马来西亚吉隆坡,玛拉理工大学医学院眼科;<sup>4</sup>马来西亚吉隆坡,马来西亚国立大学医学院妇产科;<sup>5</sup>马来西亚吉隆坡,马来西亚国立大学医学院社区卫生部门;<sup>6</sup>马来西亚吉隆坡,马来亚大学医学院眼科)

通讯作者:Tajunisah Iqbal. tajun69@yahoo.com

#### 摘要

**目的:**研究血清胰岛素样生长因子-1(IGF-1)与糖尿病性视网膜病变的联系。

**方法:**对25例怀孕的糖尿病患者及按年龄、种族、胎儿期和妊娠期相匹配的25例怀孕的非糖尿病患者的血清IGF-1水平进行测量。

**结果:**怀孕的糖尿病患者血清IGF-1水平与怀孕的非糖尿病患者血清IGF-1水平相比明显升高,24wk分别是 $366 \pm 199 \mu\text{g/L}$ 和 $184 \pm 89 \mu\text{g/L}$ ( $P=0.0001$ ),32wk分别是 $535 \pm 251 \mu\text{g/L}$ 和 $356 \pm 89 \mu\text{g/L}$ ( $P=0.007$ )。36wk分别是 $404 \pm 166 \mu\text{g/L}$ 和 $264 \pm 113 \mu\text{g/L}$ ( $P=0.003$ )。在妊娠的28,32,36wk,有明确糖尿病病史的怀孕的糖尿病患者血清IGF-1水平比妊娠期糖尿病患者血清IGF-1水平有明显的升高。在整个妊娠期怀孕的伴有视网膜病变的糖尿病患者血清IGF-1水平比那些不伴有视网膜病的患者血清IGF-1水平明显升高。

**结论:**在妊娠期提高血清IGF-1水平可能增加患糖尿病性视网膜病变的风险。

**关键词:**糖尿病性视网膜病变;怀孕;血清IGF-1水平