

Differences of bFGF gene expression in lens epithelial cells between fetuses and cataract patients

Yu-Fu Liu¹, Hong-Wei Liu^{1,2}, Yi Zhou²

¹Beijing Ophthalmology & Visual Sciences Key Lab, Tongren Eye Center, Beijing Tongren Hospital, Capital Medical University, Beijing 100005, China

²Beijing Institute of Ophthalmology, Beijing 100730, China

Correspondence to: Hong-Wei Liu. Beijing Institute of Ophthalmology, Beijing 100730, China. hw65@sohu.com

Received:2009-12-02 Accepted:2010-01-15

Abstract

• **AIM:** To study the differences of basic fibroblast growth factor (bFGF) gene expression in lens epithelial cells (LECs) between fetuses and cataract patients.

• **METHODS:** *In situ* hybridization was used to detect bFGF mRNA in the LECs that were cultured and in tissue sections from fetuses and in the LECs from the anterior capsule of cataract patients. Image analysis was used for the relative quantitative analysis of bFGF mRNA.

• **RESULTS:** bFGF gene existed in the LECs that were cultured and in tissue sections from fetuses and in the LECs from the anterior capsule of cataract patients. The integral absorbance for the fetal cultured cells, the fetal tissue sections and the capsule membrane cells of cataract patients were 627.1 ± 268.7 , 131.5 ± 42.8 and 79.2 ± 26.3 respectively. The integral absorbance of fetal cultured LECs was significantly higher than that of fetal section LECs ($P < 0.01$). The integral absorbance of cataract LECs was significantly lower than that of fetal LECs ($P < 0.01$).

• **CONCLUSION:** The *in vitro* culture of LECs can improve bFGF gene expression. The bFGF gene expression in fetal LECs is significantly higher than that in cataract LECs.

• **KEYWORDS:** lens epithelial cells; environmental scanning electron microscope; *in situ* hybridization; basic fibroblast growth factor; image analysis

DOI:10.3980/j.issn.2222-3959.2010.01.13

Liu YF, Liu HW, Zhou Y. Differences of bFGF gene expression in lens epithelial cells between fetuses and cataract patients. *Int J Ophthalmol* 2010;3(1):58–60

INTRODUCTION

Posterior capsule opacification is mainly caused by the proliferation of residual lens epithelial cells (LECs)

after operation which is caused by post operational inflammation and repair of injury, in which cytokines play an important role. The occurrence of after-cataract changes in different age groups, younger population has a higher occurrence, and children have an occurrence of 77.9%^[1]. The bFGF protein can be detected in human LECs and exists in the LECs of cataract patients as well^[2]. In this study, we compared the difference in bFGF gene expression in LECs between fetuses and cataract patients and explored the effect of cell culture on bFGF gene expression.

MATERIALS AND METHODS

Materials Reagents digoxin 1410bp bFGF cDNA probe (Academy Of Military Medical Sciences, Beijing, China), *In situ* hybridization kit and NBI/BCIP kit (Promega, Wisconsin, USA).

Methods

Preparation of specimen fetal lens capsule slides From 6 spontaneous abortuses, the left eyeballs were taken. The sclera was cut open. The anterior capsules were taken and fixed in 40g/L paraformaldehyde solution for 5 minutes and then rinsed with PBS for later use.

Cultured fetal LECs From the above 6 spontaneous abortuses, the right eyeballs were taken. The anterior capsules were cut to pieces for cell culture. Dropped the cell suspension of second generation of fetal LECs on the coverslip and cultured cells for a few days. The cells were fixed in 40g/L paraformaldehyde solution for 5 minutes and rinsed with PBS. Capsules of cataract patients: the anterior capsules from 6 cataract patients and capsules were fixed in 40g/L paraformaldehyde solution for 5 minutes and then rinsed with PBS for later use.

Examination by environmental scanning electron microscope The cultured LECs and capsules were fixed by 20g/L glutaraldehyde for 24 hours and washed by PBS for 3 times. They were examined and photographed by FEI Quanta 200 environmental scanning electron microscope.

Test on the expression of FGF gene by *in situ* hybridization Performed the experiment according to the methods described in reference^[3]. Negative control was set up. The positive cells were blue or indigo and the negative cells were colorless or slight pale red.

Image analysis Cambridge Quantimet 970 image analyzer was used for image analysis and integral absorbance of cells calculated.

Statistical Analysis Data were shown in mean \pm SD and SPSS 11 was used for *F*-test between groups.

RESULTS

bFGF Gene Expression The LECs of culture and tissue sections from fetuses expressed bFGF gene. The LECs from the anterior capsule of cataract patients expressed bFGF gene, too.

Cells Structure of ESEM The cultured cells were flat and big, the diameter was 20-22 μ m (Figure 1). The cells in capsule were prominent and small, the diameter was 5-6 μ m (Figure 2).

LEC Integral Absorbance The integral absorbance of the LECs according to image analysis on 30 cultured fetal cells, 30 tissue section cells and 30 cataract capsule cells were 627.1 \pm 268.7, 131.5 \pm 42.8 and 79.2 \pm 26.3 respectively. The integral absorbance of the cultured cells was higher than that of tissue section cells and there was great significance in the difference between the two groups ($P<0.01$). The integral absorbance of the LECs of cataract patients was lower than that of the LECs of fetuses and there was great significance between the two groups ($P<0.01$).

We also determined average absorbance which is the absorbance per unit area of cells. They were 0.031, 0.070 and 0.022 for fetal cultured cells, fetal tissue section cells and cataract capsule cells respectively. The average absorbance of fetal tissue section cells was higher than that of the cultured cells. However, since the cultured cells had a larger volume, the integral absorbance of the cultured cells was still higher than that of the tissue section cells. Both the average absorbance and the integral absorbance of the LECs of cataract patients were lower than that of the fetal cells.

DISCUSSION

There are many differences in morphology and many proteins between cultured lens epithelial cells and the lens epithelial cells of capsule^[4]. The lens is an avascular tissue, separated from the aqueous and vitreous humors by its own extracellular matrix, the lens capsule. Cell culture can injure and stimulate the tissues itself, thus can activate the synthesis of bFGF. The bFGF stimulates the proliferation of lens epithelial cells through acting on the FGF receptor in the cell membrane^[5]. The bFGF within the cell may be transported to the cell outside. However, bFGF genes can not code signal peptide sequence and how it arrives outside the cell is unclear. One of the possible causes is that the cells are dissolved. The lens capsule is also a source of




Figure 1 Cultured lens epithelial cells ($\times 500$)




Figure 2 Lens epithelial cells in capsule($\times 5000$)

bFGF, Matrix metalloproteinase 2 (MMP-2) may induce bFGF released from capsular^[6]. The bFGF can combine with extracellular matrix components such as heparin sulfate fibronectin and heparin sulfate proteoglycan. The biosynthetic genovariation of heparin sulfate can affect the binding of FGF with its receptor to reduce the effect bFGF^[7]. Some substances, such as IL1, may promote the formation of bFGF^[8].

The application of anti-sense technology can reduce bFGF mRNA expression and inhibit the promotion effect of bFGF on LECs proliferation^[9]. It is reported that the promotion effect of bFGF on LECs proliferation actualizes through MAPK/ERK signaling system^[10]. In addition, bFGF can promote FAK expression and conduct the signals of extracellular matrix into lens epithelial cells to enhance cell proliferation and migration^[11]. But, LECs proliferation was not only promoted by bFGF, but also by other factors^[12].

In situ hybridization studies on fetal tissue section, cell culture and cataract capsule have shown that there is bFGF gene expression in human lens epithelial cells, which proves that bFGF exists in these cells and plays physiological

actions when it is released to outside of cells after injury and stimulation. Posterior capsule opacification is possibly caused by injury-repair reaction after operation which induces the injured or stimulated lens epithelial cells to release bFGF. The released bFGF plus the bFGF in humor aqueous and other factors in the eye reach a certain concentration level promote cell proliferation. The capacity of the synthesis of bFGF decreases with the increase of age. The occurrence of after cataract is higher in children than in adults.

Acknowledgement: We are grateful to Dr. Xie Jiayi from Lab of Biological Imaging and Nanomedicine, National Center for Nanoscience and Technology, Beijing, China for ESEM support.

REFERENCES

- 1 Hosal BM, Biglan AW. Risk factors for secondary membrane formation after removal of pediatric cataract. *J Cataract Refract Surg* 2002;28(2):302-309
- 2 Shentu X, Yao K, Sun C, Xu W, Wu R. Expression and effect of basic fibroblast growth factor on human cataract lens epithelial cells. *Chin Med J* 2002;115(2):268-271
- 3 Liu YF, Liu HW. Expression of basic fibroblast growth factor gene in human lens epithelial cell. *Int J Ophthalmol (Guoji Yanke Zazhi)* 2008;8(7):1326-1327
- 4 Ong MD, Payne DM, Garner MH. Differential protein expression in lens epithelial whole-mounts and lens epithelial cell cultures. *Exp Eye Res* 2003;77(1):35-49
- 5 Kampmeier J, Baldysiak-Figiel A, de Jong-Hesse Y, Lang GK, Lang GE. Effect of growth factors on proliferation and expression of growth factor receptors in a human lens epithelial cell line. *J Cataract Refract Surg* 2006;32(3):510-514
- 6 Tholozan FM, Gribbon C, Li Z, Goldberg MW, Prescott AR, McKie N, Quinlan RA. FGF-2 release from the lens capsule by MMP-2 maintains lens epithelial cell viability. *Mol Biol Cell* 2007;18(11):4222-4231
- 7 Pan Y, Woodbury A, Esko JD, Grobe K, Zhang X. Heparan sulfate biosynthetic gene *Ndst1* is required for FGF signaling in early lens development. *Development* 2006;133(24):4933-4944
- 8 Sato N, Fujii A. Effects of interleukin-1alpha on the production and release of basic fibroblast growth factor in cultured nifedipine-reactive gingival fibroblasts. *J Oral Sci* 2008;50(1):83-90
- 9 Liu HW, Wang XL, Peng SL, Zhou Y, Liu DL. Inhibiting proliferation of cultured rat lens epithelial cells by bFGF antisense oligonucleotides and their liposomes. *Chin J Ophthalmol* 2004;40(8):528-532
- 10 Iyengar L, Wang Q, Rasko JE, McAvoy JW, Lovicu FJ. Duration of ERK1/2 phosphorylation induced by FGF or ocular media determines lens cell fate. *Differentiation* 2007;75(7):662-668
- 11 Kokkinos MI, Brown HJ, de Iongh RU. Focal adhesion kinase (FAK) expression and activation during lens development. *Mol Vis* 2007;13:418-430
- 12 Iyengar L, Patkunanathan B, McAvoy JW, Lovicu FJ. Growth factors involved in aqueous humour-induced lens cell proliferation. *Growth Factors* 2009;27(1):50-62